

Dr. R. Vijayaragavan

Associate Professor of Mathematics

Thiruvalluvar University

Serkkadu, Vellore-632 115.

APPLIED NUMERICAL ANALYSIS

Gaussian Elimination Method

A method to solve simultaneous linear equations
of the form $[A][X]=[C]$

Two steps

1. Forward Elimination

2. Back Substitution

Forward Elimination

The goal of forward elimination is to transform the coefficient matrix into an upper triangular matrix

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$

Forward Elimination

A set of n equations and n unknowns

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

. . . .

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = b_n$$

(n-1) steps of forward elimination

Forward Elimination

Step 1

For Equation 2, divide Equation 1 by a_{11} and multiply by a_{21}

$$\left[\frac{a_{21}}{a_{11}} \right] (a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1)$$

$$a_{21}x_1 + \frac{a_{21}}{a_{11}}a_{12}x_2 + \dots + \frac{a_{21}}{a_{11}}a_{1n}x_n = \frac{a_{21}}{a_{11}}b_1$$

Forward Elimination

Subtract the result from Equation 2.

$$\begin{array}{r} a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2 \\ - \quad a_{21}x_1 + \frac{a_{21}}{a_{11}}a_{12}x_2 + \dots + \frac{a_{21}}{a_{11}}a_{1n}x_n = \frac{a_{21}}{a_{11}}b_1 \\ \hline \left(a_{22} - \frac{a_{21}}{a_{11}}a_{12} \right)x_2 + \dots + \left(a_{2n} - \frac{a_{21}}{a_{11}}a_{1n} \right)x_n = b_2 - \frac{a_{21}}{a_{11}}b_1 \end{array}$$

or $\dot{a}_{22}x_2 + \dots + \dot{a}_{2n}x_n = \dot{b}_2$

Forward Elimination

Repeat this procedure for the remaining equations to reduce the set of equations as

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\dot{a_{32}}x_2 + \dot{a_{33}}x_3 + \dots + \dot{a_{3n}}x_n = \dot{b_3}$$

$$\begin{matrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{matrix}$$

$$\dot{a_{n2}}x_2 + \dot{a_{n3}}x_3 + \dots + \dot{a_{nn}}x_n = \dot{b_n}$$

End of Step 1

Forward Elimination

Step 2

Repeat the same procedure for the 3rd term of Equation 3.

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_{3n}}x_n = \ddot{b_3}$$

$$\begin{matrix} \cdot \\ \cdot \\ \cdot \end{matrix}$$

$$\ddot{a_{n3}}x_3 + \dots + \ddot{a_{nn}}x_n = \ddot{b_n}$$

End of Step 2

Forward Elimination

At the end of (n-1) Forward Elimination steps, the system of equations will look like

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_{3n}}x_n = \ddot{b_3}$$

.

.

.

$$a_{nn}^{(n-1)}x_n = b_n^{(n-1)}$$

End of Step (n-1)

Matrix Form at End of Forward Elimination

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a'_{22} & a'_{23} & \cdots & a'_{2n} \\ 0 & 0 & a''_{33} & \cdots & a''_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & a_{nn}^{(n-1)} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b'_2 \\ b''_3 \\ \vdots \\ b_n^{(n-1)} \end{bmatrix}$$

Back Substitution

Solve each equation starting from the last equation

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$

Example of a system of 3 equations

Back Substitution Starting Eqns

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_n}x_n = \ddot{b_3}$$

. . .

$$a_{nn}^{(n-1)}x_n = b_n^{(n-1)}$$

Back Substitution

Start with the last equation because it has only one unknown

$$x_n = \frac{b_n^{(n-1)}}{a_{nn}^{(n-1)}}$$

Back Substitution

$$x_n = \frac{b_n^{(n-1)}}{a_{nn}^{(n-1)}}$$

$$x_i = \frac{b_i^{(i-1)} - a_{i,i+1}^{(i-1)}x_{i+1} - a_{i,i+2}^{(i-1)}x_{i+2} - \dots - a_{i,n}^{(i-1)}x_n}{a_{ii}^{(i-1)}} \text{ for } i = n-1, \dots, 1$$

$$x_i = \frac{b_i^{(i-1)} - \sum_{j=i+1}^n a_{ij}^{(i-1)}x_j}{a_{ii}^{(i-1)}} \text{ for } i = n-1, \dots, 1$$

Gauss Elimination Example

Example 1

The upward velocity of a rocket is given at three different times

Table 1 Velocity vs. time data.

Time, t (s)	Velocity, v (m/s)
5	106.8
8	177.2
12	279.2



The velocity data is approximated by a polynomial as:

$$v(t) = a_1 t^2 + a_2 t + a_3, \quad 5 \leq t \leq 12.$$

Find the velocity at $t=6$ seconds .

Example 1 Cont.

Assume

$$v(t) = a_1 t^2 + a_2 t + a_3, \quad 5 \leq t \leq 12.$$

Results in a matrix template of the form:

$$\begin{bmatrix} t_1^2 & t_1 & 1 \\ t_2^2 & t_2 & 1 \\ t_3^2 & t_3 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$$

Using data from Table 1, the matrix becomes:

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Example 1 Cont.

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix} \Rightarrow \begin{bmatrix} 25 & 5 & 1 & : & 106.8 \\ 64 & 8 & 1 & : & 177.2 \\ 144 & 12 & 1 & : & 279.2 \end{bmatrix}$$

1. Forward Elimination
2. Back Substitution

Forward Elimination

Number of Steps of Forward Elimination

Number of steps of forward elimination is
 $(n-1)=(3-1)=2$

Forward Elimination: Step 1

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : 106.8 \\ 64 & 8 & 1 & : 177.2 \\ 144 & 12 & 1 & : 279.2 \end{array} \right]$$

Divide Equation 1 by 25 and multiply it by $64, \frac{64}{25} = 2.56$.

$$[25 \ 5 \ 1 \ : \ 106.8] \times 2.56 = [64 \ 12.8 \ 2.56 \ : \ 273.408]$$

Subtract the result from Equation 2

$$\begin{array}{r} [64 \ 8 \ 1 \ : \ 177.2] \\ - [64 \ 12.8 \ 2.56 \ : \ 273.408] \\ \hline [0 \ -4.8 \ -1.56 \ : \ -96.208] \end{array}$$

Substitute new equation for Equation 2

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : 106.8 \\ 0 & -4.8 & -1.56 & : -96.208 \\ 144 & 12 & 1 & : 279.2 \end{array} \right]$$

Forward Elimination: Step 1 (cont.)

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : & 106.8 \\ 0 & -4.8 & -1.56 & : & -96.208 \\ 144 & 12 & 1 & : & 279.2 \end{array} \right] \begin{matrix} \text{Divide Equation 1 by 25 and} \\ \text{multiply it by 144, } \frac{144}{25} = 5.76. \end{matrix}$$

$$[25 \ 5 \ 1 \ : \ 106.8] \times 5.76 = [144 \ 28.8 \ 5.76 \ : \ 615.168]$$

Subtract the result from
Equation 3

$$\begin{array}{r} [144 \ 12 \ 1 \ : \ 279.2] \\ - [144 \ 28.8 \ 5.76 \ : \ 615.168] \\ \hline [0 \ -16.8 \ -4.76 \ : \ -335.968] \end{array}$$

Substitute new equation
for Equation 3

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : & 106.8 \\ 0 & -4.8 & -1.56 & : & -96.208 \\ 0 & -16.8 & -4.76 & : & -335.968 \end{array} \right]$$

Forward Elimination: Step 2

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : & 106.8 \\ 0 & -4.8 & -1.56 & : & -96.208 \\ 0 & -16.8 & -4.76 & : & -335.968 \end{array} \right] \quad \begin{array}{l} \text{Divide Equation 2 by } -4.8 \\ \text{and multiply it by } -16.8, \\ \frac{-16.8}{-4.8} = 3.5 \end{array}$$

$$[0 \quad -4.8 \quad -1.56 \quad : \quad -96.208] \times 3.5 = [0 \quad -16.8 \quad -5.46 \quad : \quad -336.728]$$

Subtract the result from
Equation 3

$$\begin{array}{r} [0 \quad -16.8 \quad -4.76 \quad : \quad 335.968] \\ - [0 \quad -16.8 \quad -5.46 \quad : \quad -336.728] \\ \hline [0 \quad 0 \quad 0.7 \quad : \quad 0.76] \end{array}$$

Substitute new equation for
Equation 3

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : & 106.8 \\ 0 & -4.8 & -1.56 & : & -96.208 \\ 0 & 0 & 0.7 & : & 0.76 \end{array} \right]$$

Back Substitution

Back Substitution

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : 106.8 \\ 0 & -4.8 & -1.56 & : -96.2 \\ 0 & 0 & 0.7 & : 0.7 \end{array} \right] \Rightarrow \left[\begin{array}{ccc|c} 25 & 5 & 1 & : 106.8 \\ 0 & -4.8 & -1.56 & : -96.2 \\ 0 & 0 & 0.7 & : 0.7 \end{array} \right] \left[\begin{array}{c} a_1 \\ a_2 \\ a_3 \end{array} \right] = \left[\begin{array}{c} 106.8 \\ -96.2 \\ 0.76 \end{array} \right]$$

Solving for a_3

$$0.7a_3 = 0.76$$

$$a_3 = \frac{0.76}{0.7}$$

$$a_3 = 1.08571$$

Back Substitution (cont.)

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.208 \\ 0.76 \end{bmatrix}$$

Solving for a_2

$$-4.8a_2 - 1.56a_3 = -96.208$$

$$a_2 = \frac{-96.208 + 1.56a_3}{-4.8}$$

$$a_2 = \frac{-96.208 + 1.56 \times 1.08571}{-4.8}$$

$$a_2 = 19.6905$$

Back Substitution (cont.)

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.2 \\ 0.76 \end{bmatrix}$$

Solving for a_1

$$25a_1 + 5a_2 + a_3 = 106.8$$

$$\begin{aligned} a_1 &= \frac{106.8 - 5a_2 - a_3}{25} \\ &= \frac{106.8 - 5 \times 19.6905 - 1.08571}{25} \\ &= 0.290472 \end{aligned}$$

Gaussian Elimination Solution

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 0.290472 \\ 19.6905 \\ 1.08571 \end{bmatrix}$$

Example 1 Cont.

Solution

The solution vector is

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 0.290472 \\ 19.6905 \\ 1.08571 \end{bmatrix}$$

The polynomial that passes through the three data points is then:

$$\begin{aligned} v(t) &= a_1 t^2 + a_2 t + a_3 \\ &= 0.290472 t^2 + 19.6905 t + 1.08571, \quad 5 \leq t \leq 12 \end{aligned}$$

$$\begin{aligned} v(6) &= 0.290472(6)^2 + 19.6905(6) + 1.08571 \\ &= 129.686 \text{ m/s.} \end{aligned}$$

Gauss-Seidel Method

Gauss-Seidel Method

An iterative method.

Basic Procedure:

- Algebraically solve each linear equation for x_i
- Assume an initial guess solution array
- Solve for each x_i and repeat
- Use absolute relative approximate error after each iteration to check if error is within a pre-specified tolerance.

Gauss-Seidel Method

Why?

The Gauss-Seidel Method allows the user to control round-off error.

Elimination methods such as Gaussian Elimination and LU Decomposition are prone to round-off error.

Also: If the physics of the problem are understood, a close initial guess can be made, decreasing the number of iterations needed.

Gauss-Seidel Method

Algorithm

A set of n equations and n unknowns:

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

$$\begin{matrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{matrix}$$

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = b_n$$

If: the diagonal elements are non-zero

Rewrite each equation solving for the corresponding unknown

ex:

First equation, solve for x_1

Second equation, solve for x_2

Gauss-Seidel Method

Algorithm

Rewriting each equation

$$x_1 = \frac{c_1 - a_{12}x_2 - a_{13}x_3 - \dots - a_{1n}x_n}{a_{11}} \quad \longleftarrow \text{From Equation 1}$$

$$x_2 = \frac{c_2 - a_{21}x_1 - a_{23}x_3 - \dots - a_{2n}x_n}{a_{22}} \quad \longleftarrow \text{From equation 2}$$
$$\vdots \quad \vdots \quad \vdots$$

$$x_{n-1} = \frac{c_{n-1} - a_{n-1,1}x_1 - a_{n-1,2}x_2 - \dots - a_{n-1,n-2}x_{n-2} - a_{n-1,n}x_n}{a_{n-1,n-1}} \quad \longleftarrow \text{From equation n-1}$$

$$x_n = \frac{c_n - a_{n1}x_1 - a_{n2}x_2 - \dots - a_{n,n-1}x_{n-1}}{a_{nn}} \quad \longleftarrow \text{From equation n}$$

Gauss-Seidel Method

Algorithm

General Form of each equation

$$x_1 = \frac{c_1 - \sum_{\substack{j=1 \\ j \neq 1}}^n a_{1j} x_j}{a_{11}}$$
$$x_{n-1} = \frac{c_{n-1} - \sum_{\substack{j=1 \\ j \neq n-1}}^n a_{n-1,j} x_j}{a_{n-1,n-1}}$$
$$x_2 = \frac{c_2 - \sum_{\substack{j=1 \\ j \neq 2}}^n a_{2j} x_j}{a_{22}}$$
$$x_n = \frac{c_n - \sum_{\substack{j=1 \\ j \neq n}}^n a_{nj} x_j}{a_{nn}}$$

Gauss-Seidel Method

Algorithm

General Form for any row ‘i’

$$x_i = \frac{c_i - \sum_{\substack{j=1 \\ j \neq i}}^n a_{ij} x_j}{a_{ii}}, i = 1, 2, \dots, n.$$

How or where can this equation be used?

Gauss-Seidel Method

Solve for the unknowns

Assume an initial guess for [X]

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_{n-1} \\ x_n \end{bmatrix}$$

Use rewritten equations to solve for each value of x_i .

Important: Remember to use the most recent value of x_i . Which means to apply values calculated to the calculations remaining in the **current** iteration.

Gauss-Seidel Method

Calculate the Absolute Relative Approximate Error

$$|\epsilon_a|_i = \left| \frac{x_i^{new} - x_i^{old}}{x_i^{new}} \right| \times 100$$

So when has the answer been found?

The iterations are stopped when the absolute relative approximate error is less than a prespecified tolerance for all unknowns.

Gauss-Seidel Method: Example 1

The upward velocity of a rocket is given at three different times

Table 1 Velocity vs. Time data.

Time, t (s)	Velocity v (m/s)
5	106.8
8	177.2
12	279.2



The velocity data is approximated by a polynomial as:

$$v(t) = a_1 t^2 + a_2 t + a_3, \quad 5 \leq t \leq 12.$$

Gauss-Seidel Method: Example 1

Using a Matrix template of the form

$$\begin{bmatrix} t_1^2 & t_1 & 1 \\ t_2^2 & t_2 & 1 \\ t_3^2 & t_3 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix}$$

The system of equations becomes

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Initial Guess: Assume an initial guess of

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$$

Gauss-Seidel Method: Example 1

Rewriting each equation

$$a_1 = \frac{106.8 - 5a_2 - a_3}{25}$$

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

$$a_2 = \frac{177.2 - 64a_1 - a_3}{8}$$

$$a_3 = \frac{279.2 - 144a_1 - 12a_2}{1}$$

Gauss-Seidel Method: Example 1

Applying the initial guess and solving for a_i

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix}$$

Initial Guess

$$a_1 = \frac{106.8 - 5(2) - (5)}{25} = 3.6720$$

$$a_2 = \frac{177.2 - 64(3.6720) - (5)}{8} = -7.8510$$

$$a_3 = \frac{279.2 - 144(3.6720) - 12(-7.8510)}{1} = -155.36$$

When solving for a_2 , how many of the initial guess values were used?

Gauss-Seidel Method: Example 1

Finding the absolute relative approximate error

$$|\epsilon_a|_i = \left| \frac{x_i^{new} - x_i^{old}}{x_i^{new}} \right| \times 100$$

$$|\epsilon_a|_1 = \left| \frac{3.6720 - 1.0000}{3.6720} \right| \times 100 = 72.76\%$$

$$|\epsilon_a|_2 = \left| \frac{-7.8510 - 2.0000}{-7.8510} \right| \times 100 = 125.47\%$$

$$|\epsilon_a|_3 = \left| \frac{-155.36 - 5.0000}{-155.36} \right| \times 100 = 103.22\%$$

At the end of the first iteration

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 3.6720 \\ -7.8510 \\ -155.36 \end{bmatrix}$$

The maximum absolute relative approximate error is 125.47%

Gauss-Seidel Method: Example 1

Iteration #2

Using

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 3.6720 \\ -7.8510 \\ -155.36 \end{bmatrix}$$

from iteration #1

the values of a_i are found:

$$a_1 = \frac{106.8 - 5(-7.8510) - 155.36}{25} = 12.056$$

$$a_2 = \frac{177.2 - 64(12.056) - 155.36}{8} = -54.882$$

$$a_3 = \frac{279.2 - 144(12.056) - 12(-54.882)}{1} = -798.34$$

Gauss-Seidel Method: Example 1

Finding the absolute relative approximate error

$$|\epsilon_a|_1 = \left| \frac{12.056 - 3.6720}{12.056} \right| \times 100 = 69.543\%$$

$$|\epsilon_a|_2 = \left| \frac{-54.882 - (-7.8510)}{-54.882} \right| \times 100 = 85.695\%$$

$$|\epsilon_a|_3 = \left| \frac{-798.34 - (-155.36)}{-798.34} \right| \times 100 = 80.540\%$$

At the end of the second iteration

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 12.056 \\ -54.882 \\ -798.54 \end{bmatrix}$$

The maximum absolute relative approximate error is 85.695%

Gauss-Seidel Method: Example 1

Repeating more iterations, the following values are obtained

Iteration	a_1	$ e_a _1 \%$	a_2	$ e_a _2 \%$	a_3	$ e_a _3 \%$
1	3.6720	72.767	-7.8510	125.47	-155.36	103.22
2	12.056	69.543	-54.882	85.695	-798.34	80.540
3	47.182	74.447	-255.51	78.521	-3448.9	76.852
4	193.33	75.595	-1093.4	76.632	-14440	76.116
5	800.53	75.850	-4577.2	76.112	-60072	75.963
6	3322.6	75.906	-19049	75.972	-249580	75.931

Notice – The relative errors are not decreasing at any significant rate

Also, the solution is not converging to the true solution of $\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 0.29048 \\ 19.690 \\ 1.0857 \end{bmatrix}$

THE END